

CHARACTERISTIC COHOMOLOGY OF MUMFORD-TATE DOMAINS

MOHAMMAD REZA RAHMATI, KHURRAM SHABBIR

ABSTRACT. In this text we pretend to find generators and relations for the universal characteristic cohomology of MT-domains of Hodge structures by some structure theorem via toric stacks. We also compare a uniformization theorem of toric stacks with the corresponding structure on MT-domains.

INTRODUCTION

Assume D is a period domain of pure Hodge structures defined by Griffiths, with the period map $(\Delta^*)^n \rightarrow \Gamma \backslash D$, where Γ is the monodromy group. The domain D can be understood as the orbit of a polarized Hodge structure $(H_{\mathbb{Q}}, F^{\bullet}, Q)$, under the action of a reductive group G preserving the polarization form Q . We treat a partial compactification of $\Gamma \backslash D$ so that the period map is extended over Δ^n . There are different theories of compactifications for period domains D . Kato and Usui provide the toroidal compactifications for any period domain D , which is not Hermitian symmetric in general and show these are moduli spaces of log Hodge structures. This partial compactification is given by using toroidal embedding associated to the cone generated by the data of the monodromies. The Mumford-Tate (MT)-domains of a Hodge structure

$$\phi : \mathbb{S}(\mathbb{R}) \rightarrow Gl(V, \mathbb{R})$$

is the orbit under the MT-group of the Hodge structure ϕ . They are reductive Lie algebraic group defined over $\mathbb{Q}$.

Mumford-Tate domains can be considered as a toric stack, by a definition using Stacky-fans. The structure is compatible with the Kato-Usui generalized toroidal compactification as we will see below by the definition. In this way we may regard some additional data on MT-domains which preserves the stabilizers of the points under a Lie group action. A toric stack is a simple generalization of the definition of toric variety, but we will concern the toric action as a groupoid action. There are several definitions of toric Stacks in the literature. A toric stack is the stack quotient

1991 *Mathematics Subject Classification.* 14D23.

Key words and phrases. Deligne-Mumford Stacks, Toric variety, Mumford-Tate domains, Gerbe, Root Stack, Stacky fan, Cohomology of toric stacks.

of a normal toric variety X , by a subgroup G of its Torus T , cf. [GS]. The main task of this text is to match this definition with how to express the definition with that of period domains of Griffiths, Mumford-Tate domains, and their boundary points via the Kato-Usui generalized toroidal compactification. As an application one can consider the universal characteristic cohomology of Mumford-Tate domains as a type of stack cohomology (orbifold cohomology).

A simple way to define a toric stack is by the definition of a stacky fan (Σ, β, N) , where Σ is a fan, $\beta : \mathbb{Z}^n \rightarrow N$ a homomorphism of abelian groups with finite cokernel, and N is a lattice in $\mathbb{R}^n$. Toric stacks are Deligne-Mumford stacks, that is they can be understood similar to usual manifolds. Such stacks are also smooth as DM stacks and their boundary points have codimension 1 as a sub-stack. Toric stacks satisfy a simple uniformization structures as a global quotient via the structure of their Chow group or equivalently their Picard group of line bundles. We explain this by root sheaves or root of line bundles associated to rays in their fan.

In this text we show that MT-domains of Hodge structure are smooth DM-stacks and in this way satisfy the same uniformization theorem as general smooth DM-stacks. We propose to apply the structure theorems of toric stacks to Kato-Usui partial toric compactification of period and Mumford-Tate domains. To this target, we briefly mention some basic results on toric stacks as smooth Deligne-Mumford stacks from the references, [FMN] and [J]. Our strategy is to combine this literature with Hodge theory as much as possible. The Proposition 2.1 and part (4) of Section 3 seem to have new contributions. The basic material on MT-domains in use are from [TH], [KP], [GGK]. We refer to [WF] for preliminaries on Toric varieties.

1. TORIC STACKS

Toric stacks are generalization of the notion of toric varieties and their torus actions. The point is to upgrade the action of the torus to a groupoid action. The action ρ is equivalently thought of a functor $\rho : BG \rightarrow \mathbf{Sets}$, from the group G regarded as a one-object groupoid, denoted by BG . This functor sends the single object to the set X . The notion of groupoids and their action can be generalized over arbitrary schemes. Let S be a base scheme. The action of the groupoid $\mathcal{G}$ on an S -stack is a morphism $\mathcal{G} \times_S X \rightarrow X$ with two arrows $\eta_x : e \times x \Rightarrow x$, $(g_1.g_2) \times x \Rightarrow g_1.(g_2.x)$. The example we are interested in are Deligne-Mumford torus. A Deligne-Mumford torus is a Picard stack over $\mathrm{Spec}(\mathbb{C})$ which is obtained as a quotient $[T_L/G]$, with $\phi : L \rightarrow N$ is a morphism of finitely generated abelian groups such that $\ker(\phi)$ is free and $\mathrm{coker}(\phi) =: G$ is finite, and T_L is a torus associated to the lattice L . Any Deligne-Mumford (DM) torus is isomorphic as Picard stack to $T \times BG$, where T is a torus and G is a finite abelian group.

A smooth toric Deligne-Mumford stack is a smooth separated DM-stack X together with an open immersion of a Deligne-Mumford torus $\iota : \mathcal{T} \hookrightarrow X$ with dense image such that the action of $\mathcal{T}$ on itself extends to an action $\mathcal{T} \times X \rightarrow X$. In this case a morphism is a morphism of stacks which extends a morphism of Deligne-Mumford tori. A toric orbifold is a toric DM-stack with generically trivial stabilizer. A toric DM-stack is a toric orbifold iff its DM-torus is an ordinary torus. The same way to define toric varieties by combinatorial data of their fan, a toric stack may also be defined by a stacky fan.

A stacky fan is a pair (Σ, β) , where Σ is a fan on a lattice L and $\beta : L \rightarrow N$ is a homomorphism to a lattice N , so that $\text{coker}(\beta)$ is finite. A stacky fan gives rise to a toric stack as follows. Let X_Σ be the toric variety associated to Σ . The map $\beta^* : N^* \rightarrow L^*$ induces a homomorphism of Tori, $T_\beta : T_L \rightarrow T_N$, by naturally identifying β with the induced map on lattices of 1-parameter subgroups. Since $\text{coker}(\beta)$ is finite, β^* is injective, so T_β is surjective. Let $G_\beta = \ker T_\beta$. Note that T_L is the torus on X_Σ , and $G_\beta \subset T_L$ is a subgroup. The action of G_β on X_Σ is induced by the homomorphism $G_\beta \rightarrow T_L$. If (Σ, β) is a stacky fan, we define the toric stack $\mathcal{X}_{\Sigma, \beta}$ to be $[\mathcal{X}/G_\beta]$, with the torus $T_N = T_L/G_\beta$. This construction is also reversible in a trivial way, [GS].

One may expect that toric DM-stack can be represented by a larger Z with larger group G . For instance, the classifying stack $B\mu_3 = [*/\mu_3]$ is a toric DM stack by the stacky fan $(N = \mu_3, 0, 0)$. It is isomorphic to $[\mathbb{C}^\times/\mathbb{C}^\times]$ where $\mathbb{C}^\times$ acts by $(\lambda, x) \rightarrow \lambda^3 x$. It is simply possible that a toric stack correspond to different stacky fans. Motivated by the study of gerbes one introduces the notion of extended stacky fan. An extended stacky fan is a triple (N, Σ, β^e) where N, Σ are the same as the stacky fan Σ , and $\beta^e : \mathbb{Z}^m \rightarrow N$ is determined by $b_1, \dots, b_n$ and additional elements $b_{n+1}, \dots, b_m$. Then the toric DM-stack is defined by $[Z^e/G^e]$, where $Z^e = Z \times (\mathbb{C}^\times)^{m-n}$, and G^e acts on Z^e through the homomorphism $\alpha^e : G^e \rightarrow (\mathbb{C}^\times)^{m-n}$ determined by the extended stacky fan, [J].

A coherent sheaf on a DM stack $[Z/G]$ is a G -equivariant sheaf on Z . As usual a line bundle on $[Z/G]$ is given by a character χ of G . Let b be a positive integer. We denote by $\sqrt[b]{L/X}$ the fiber product

$$(1) \quad \begin{array}{ccc} \sqrt[b]{L/X} & \longrightarrow & B\mathbb{C}^* \\ \downarrow & & \downarrow \wedge b \\ X & \xrightarrow{L} & B\mathbb{C}^* \end{array}$$

where $\wedge b$ is the map that sends an invertible sheaf to its b -th power. More explicitly an object of $\sqrt[b]{L/X}$ over $f : S \rightarrow X$ is a couple (M, ϕ) where M is an invertible sheaf on S and $\phi : M^{\otimes b} \xrightarrow{\cong} f^*L$ is an isomorphism. Let X be a smooth algebraic stack.

Assume $D = (D_1, \dots, D_n)$ are n effective Cartier divisors and $a = (a_1, \dots, a_n) \in \mathbb{N}_{>0}^n$. Then the a -th root of (X, D) is the fiber product

$$(2) \quad \begin{array}{ccc} \sqrt[a]{D/X} & \longrightarrow & [\mathbb{A}^n/(\mathbb{C}^*)^n] \\ \downarrow & & \downarrow \wedge a \\ X & \xrightarrow{L} & [\mathbb{A}^n/(\mathbb{C}^*)^n] \end{array}$$

$\wedge a : [\mathbb{A}^n/(\mathbb{C}^*)^n] \rightarrow [\mathbb{A}^n/(\mathbb{C}^*)^n]$ is the stack morphism $x_i \rightarrow x_i^{a_i}$ and $\lambda_i \rightarrow \lambda_i^{a_i}$, where x_i, λ_i are coordinates in $\mathbb{A}^n$ and $(\mathbb{C}^*)^n$ respectively. The structure morphism $X \rightarrow \bar{X}$ into its coarse moduli space factors canonically via the toric morphisms $X \rightarrow X^{\text{rig}} \rightarrow X^{\text{can}} \rightarrow \bar{X}$ where the first map is an abelian gerbe over X^{rig} , and is a fiber product of roots of toric divisors as in (2), and the last is the minimal orbifold having $\bar{X}$ as a coarse moduli space. For instance, the stack $\mathbb{P}(w) \cong [(\mathbb{C}^{n+1} \setminus 0)/\mathbb{C}^*]$ is a toric DM-stack with deligne-Mumford torus $[\mathbb{C}^{n+1}/\mathbb{C}^*] \cong \mathbb{C}^n \times B\mu_d$. It is canonical if $\gcd(w_0, \dots, \hat{w}_i, \dots, w_n) = 1$. It is an orbifold if $\gcd(w_0, \dots, w_n) = 1$, [FMN].

Proposition 1.1. [FMN] (1) Let X be toric Deligne-Mumford stack with Deligne-Mumford torus isomorphic to $T \times BG$. Given $G = \prod_{j=1}^l \mu_{b_j}$. There exists L_j in $\text{Pic}(X^{\text{rig}})$ such that X is isomorphic as a G -gerbe over X^{rig} to

$$(3) \quad \sqrt[b_1]{L_1/X^{\text{rig}}} \times_{X^{\text{rig}}} \dots \times_{X^{\text{rig}}} \sqrt[b_l]{L_l/X^{\text{rig}}}$$

where the classes $(L_1, \dots, L_l) \in \prod_{j=1}^l \text{Pic}(X^{\text{rig}})/b_j \text{Pic}(X^{\text{rig}})$ are unique. Moreover the reduced closed substack $X \setminus \mathcal{T}$ is a simple normal crossing divisor.

(2) Let X be a smooth toric Deligne-Mumford stack with DM-torus isomorphic to $T \times BG$, with $\bar{X}$ as coarse moduli. denote by n the number of rays in the fan of $\bar{X}$. Then there exists a unique $(a_1, \dots, a_n) \in \mathbb{N}_{>}^n$ such that

$$X^{\text{rig}} \cong \sqrt[a_1]{D_1^{\text{can}}/X^{\text{can}}} \times_{X^{\text{can}}} \dots \times_{X^{\text{can}}} \sqrt[a_n]{D_n^{\text{can}}/X^{\text{can}}}$$

where D_i is the divisor corresponding to the ray ρ_i .

2. HIGHER STRUCTURE ON MUMFORD-TATE DOMAINS

We apply the structure theorems of section 1 to the structure on period and Mumford-Tate domains. To begin with let V be finite dimensional $\mathbb{Q}$ -vector space, and Q a non-degenerate bilinear map $Q : V \otimes V \rightarrow \mathbb{Q}$ which is $(-1)^n$ -symmetric for some fixed n . A Hodge structure is given by a representation

$$\phi : \mathbb{U}(\mathbb{R}) \rightarrow \text{Aut}(V, Q)_{\mathbb{R}}, \quad \mathbb{U}(\mathbb{R}) = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}, \quad a^2 + b^2 = 1$$

It decomposes over $\mathbb{C}$ into eigenspaces $V^{p,q}$ such that $\phi(t).u = t^p \bar{t}^q .u$ for $u \in V^{p,q}$, and $\overline{V^{p,q}} = V^{q,p}$. It is well-known that $\text{Ad}(\phi) : \mathbb{U}(\mathbb{R}) \rightarrow \text{Aut}(\mathfrak{g}_{\mathbb{R}}, B)$ defines a weight 0 Hodge structure, where B is the Killing form. Then $\mathfrak{g}_{\mathbb{C}}$ has a decomposition

$$\mathfrak{g} = \mathfrak{g}_{\phi}^{-} \oplus \overline{\mathfrak{g}_{\phi}^{-}} \oplus \mathfrak{h}_{\phi}, \quad \mathfrak{g}^{-} := \oplus_{i>0} \mathfrak{g}_{\phi}^{-i,i}$$

and $T_{\phi, \mathbb{R}} D \otimes \mathbb{C} = \mathfrak{g}_{\phi}^{-} \oplus \overline{\mathfrak{g}_{\phi}^{-}}$, $T_{\phi} D = \mathfrak{g}_{\phi}^{-}$. The Mumford-Tate group M_{Φ} of the Hodge structure ϕ is the smallest $\mathbb{Q}$ -algebraic subgroup containing the image of ϕ , and is a reductive Lie group, that satisfies the almost product decomposition

$$(4) \quad M_{\Phi} = M_1 \times \dots \times M_l \times T$$

The associated Mumford-Tate domains D_i (the $M_i(\mathbb{R})$ -orbit of $\phi(s_0)$) have a splitting of the

$$(5) \quad D_{M_{\Phi}} = D_1 \times \dots \times D_l$$

Assume $\Phi : S \rightarrow \Gamma \setminus D$ is a variation of Hodge structure and let $T_1, \dots, T_n$ be generators of the monodromy group Γ . Then the partial compactification $\Gamma \setminus D_{\sigma}$ is given by the cone

$$\sigma = \sum_{j=1}^n \mathbb{R}_{\geq 0} N_j, \quad N_j = \log T_j$$

in the lie algebra $\mathfrak{g} := \text{Lie}(G_{\mathbb{C}}) \subset \text{Hom}(V, V)$. Here a boundary point is a nilpotent orbit associated to a face σ . Set $\Gamma(\sigma)^{\text{gp}} = \exp(\sigma_{\mathbb{R}}) \cap G_{\mathbb{Z}}$, $\Gamma(\sigma) = \exp(\sigma) \cap G_{\mathbb{Z}}$. The monoid $\Gamma(\sigma)$ defines the toric variety

$$(6) \quad D_{\sigma} := \text{Spec}([\mathbb{C}[\Gamma(\sigma)^{\vee}]]_{an} \cong \text{Hom}(\Gamma(\sigma)^{\vee}, \mathbb{C})$$

with the torus

$$(7) \quad T_{\sigma} := \text{Spec}(\mathbb{C}[\Gamma(\sigma)^{\vee \text{gp}}])_{an} \cong \text{Hom}(\Gamma(\sigma)^{\vee \text{gp}}, \mathbb{G}_m) \cong \mathbb{G}_m \otimes \Gamma(\sigma)^{\text{gp}}$$

Let Σ be a maximal fan in $\mathfrak{m} := \text{Lie}(M)$. Consider the injective map $\beta = e : \log \gamma \mapsto \gamma$ restricted to $\Sigma = \mathbb{Z}\langle N_1, \dots, N_r \rangle$. The map e is an isomorphism onto its image $\mathfrak{m}_{\mathbb{Z}}$ a maximal lattice in $\mathfrak{m}$. The triple $(D_M, e : \Sigma_{\mathbb{Z}} \rightarrow \mathfrak{m}_{\mathbb{Z}}, \Sigma)$ is a toric stack. On the lattices e is an isomorphism, mentioning that the generic stabilizer is trivial.

Theorem 2.1. *The triple $(D_M, e : \Sigma_{\mathbb{Z}} \rightarrow \mathfrak{m}_{\mathbb{Z}}, \Sigma)$ is a smooth toric Deligne-Mumford stack. This stacky structure decomposes as*

$$(8) \quad D_{M_{\Phi}} = D_1 \times \dots \times D_l$$

where

$$\begin{aligned} D &\cong \sqrt[b_1]{L_1/D} \times_D \dots \times_D \sqrt[b_l]{L_l/D} \cong \sqrt[d]{L/Z} \\ D_i &\cong \sqrt[b_i]{L_{1i}/D_i} \times_{D_i} \dots \times_{D_i} \sqrt[b_{li}]{L_{li}/D_i} \cong \sqrt[d_i]{L_i/Z_i} \end{aligned}$$

where $b_{ij} \in \mathbb{N}_{>0}$ and $d = \gcd(b_1, \dots, b_l)$, $d_i = \gcd(b_{1i}, \dots, b_{li})$. Moreover $D_{M_{\Phi}}$ is rigid.

Proof. The first part is by the definition and the construction above. The decomposition in (8) follows from proposition 1.1 and the uniqueness of splitting in (5). It is rigid because the map $e : \Sigma_{\mathbb{Z}} \rightarrow \mathfrak{m}_{\mathbb{Z}}$ is an isomorphism onto its image. $\square$

3. COHOMOLOGY

(1) Differentiable stacks are stacks over the category of differentiable manifolds. They are the stacks associated to Lie groupoids. A groupoid $X_1 \rightrightarrows X_0$ is a Lie groupoid if X_0 and X_1 are differentiable manifolds, structure maps are differentiable, source and target maps are submersion. There is associated a simplicial nerve X_p of p -fold cartesian product of X_1 over X_0 . Then we get an associated co-simplicial object $(\Omega^q(X^\bullet, \partial = \sum_{i=0}^p (-1)^i \partial_i^*))$. The cohomology groups $H^k(X, \Omega^q)$ are called the Cech cohomology groups of the groupoid $X = [X_1 \rightrightarrows X_0]$. The double complex $A^{pq} := \Omega^q(X_p)$ is called the de Rham complex of the stack $\mathcal{X}$, and its cohomologies are called de Rham cohomologies of $\mathcal{X}$. The cohomology of the DM-stack X is defined to be $H_{DR}^n(X) = H_{DR}^n(X_1 \rightrightarrows X_0)$, for any groupoid atlas $X_1 \rightrightarrows X_0$ of the stack X .

If G is a Lie group then $H^k(BG, \Omega^0)$, is the group cohomology of G calculated with differentiable cochains. When the stack X is a quotient stack there is a clear explanation of its cohomology as an equivariant cohomology. There is a well-known generalization of the de Rham complex to the equivariant case, namely the Cartan complex $\Omega_G^\bullet(X)$, defined by

$$(9) \quad \Omega_G^\bullet(X) := \bigoplus_{2k+i=n} (S^k \mathfrak{g}^\vee \otimes \Omega^i(X))^G$$

where $S^\bullet \mathfrak{g}^\vee$ is the symmetric algebra on the dual of the Lie algebra of G . The group G acts on $\mathfrak{g}$ by adjoint representation on $\mathfrak{g}^\vee$ and by pull back of differential forms

on $\Omega^\bullet(X)$. The Cartan differential is $d_{DR} - \iota$ where ι is the tensor induced by the vector bundle homomorphism $\mathfrak{g}_X \rightarrow T_X$ coming from differentiating the action. If G is compact, the augmentation is a quasi-isomorphism, i.e

$$(10) \quad H_G^i(X) \xrightarrow{\cong} H^i(Tot \Omega_G^\bullet(X_\bullet))$$

for all i and the groupoid $X_\bullet$, [KB].

Proposition 3.1. [KB] *If the Lie group G is compact, there is a natural isomorphism*

$$H_G^i(X) \xrightarrow{\cong} H_{DR}^i(G \times X \rightrightarrows X) = H_{DR}^i([X/G])$$

As a corollary $H_{DR}^*(BG) = (S^{2*}\mathfrak{g}^\vee)^G$ for a compact lie group G . If G is not compact, then $H_{DR}([X/G])$ is still equal to equivariant cohomology. This fact holds for equivariant cohomology in general.

(2) Let X be a complex manifold and $W \subset X$ a holomorphic sub-bundle. Then $I = W^\perp \subset T^*(X)$ is also holomorphic and let $\bar{I}$ be its conjugate. Let $\mathcal{I}^{\bullet\bullet} \subset \mathcal{A}^{\bullet\bullet}$ be the differential ideal of sections of $I \oplus \bar{I}$ in the algebra of smooth differential forms on X . The characteristic cohomology of X denoted $H_X^*(X)$ is the cohomology of the double complex $\subset \mathcal{A}^{\bullet\bullet}/\mathcal{I}^{\bullet\bullet}$.

Let D be a period domain for a PHS (V, Q, ϕ) of weight n and set $\Gamma_{\mathbb{Z}} = Aut(V_{\mathbb{Z}}, \mathbb{Q})$. In the tangent bundle TD there is a homogeneous sub-bundle W whose fiber at ϕ is $W_\phi = \mathfrak{g}_\phi^{-1,1} = \{\psi \in T_\phi D : \psi(F_\phi^p) \subset F_\phi^{p+1}\}$ defined by, namely infinitesimal period relations (IPR). There is a natural inclusion $T_{F^\bullet} \check{D} \subset \oplus_p Hom(F^p, V_{\mathbb{C}}/F^p)$. We shall assume that, the canonical sub-bundle $W \subset T\check{D}$ given by the infinitesimal period relation is defined by

$$W_{F^\bullet} = T_{F^\bullet} \check{D} \cap (\oplus_p Hom(F^p, F^{p-1}/F^p))$$

The bundle $W \rightarrow \check{D}$ is acted by $G_{\mathbb{C}}$, and the action of $G_{\mathbb{R}}$ on $W \rightarrow D$ leaves invariant the metric given by Cartan killing form at each point. With the identification $T_\phi D = \oplus_{i>0} \mathfrak{g}^{-i,i}$ we have $W_\phi = \mathfrak{g}_\phi^{-1,1}$. In our case, take $X = D_M \subset D$ to be a Mumford-Tate domain, and $W_M \subset TD_M$ is the infinitesimal period relation. Denote by $\Lambda_M^{\bullet\bullet}$ the complex of $G(\mathbb{R})$ -invariant forms in $A^{\bullet\bullet}(D_M)/\mathcal{I}^{\bullet\bullet}$ with the operator $\delta : \Lambda_M^\bullet \rightarrow \Lambda_M^{\bullet+1}$. $H^*(\Lambda_M^\bullet, \delta_M)$ is called the universal characteristic cohomology, [GGK].

Proposition 3.2. [GGK] *The universal characteristic cohomology i.e the cohomology of $A^{\bullet\bullet}(D_M)/\mathcal{I}^{\bullet\bullet}, \delta : \Lambda_M^\bullet \rightarrow \Lambda_M^{\bullet+1} := H^*(\Lambda_M^\bullet, \delta_M)$ is equal to the equivariant cohomology $H^*(I^{\bullet\bullet})$. Moreover, $H^{2p-1}(\Lambda_M^\bullet, \delta_M) = 0$ and $H^{2p-1}(\Lambda_M^\bullet, \delta_M) = (\Lambda^{p,p})^{m^{0,0}}$.*

The universal characteristic cohomology is equal to the de Rham cohomology of the stack D_M . In case $D_M = D$ and $M = G$, the universal characteristic cohomology

is generated by the chern forms of Hodge bundles. According to the proposition 3.2 and the remark after that, the universal characteristic cohomology can be understood as the stack cohomology of the Mumford-Tate domains. One can ask; hat are the conditions that the chern forms of Hodge bundles on D_M generate the characteristic cohomology? In case this holds what are the relations?.

(3) Keeping the notation as section (1), let $P \rightarrow B$ be a principal $(\mathbb{C}^\times)^m$ -bundle. The group G^e acts on the fiber product $P \times_{(\mathbb{C}^\times)^m} Z^e$ via the map α^e , where α^e is given by the stacky fan. Each $\theta \in M = N^*$, gives $\chi_\theta : (\mathbb{C}^\times)^m \rightarrow (\mathbb{C}^\times)$ and assume the line bundle $\psi_\theta = P \times_{\chi_\theta} \mathbb{C}$ be given by the ray θ . Then one knows that

$$(11) \quad H_{orb}^*([(P \times_{\alpha^e} Z^e)/G^e]) = \frac{H^*(B)[N] \otimes \mathbb{Q}[N]^{\Sigma^e}}{\langle \sum_i c_1(\psi_\theta) + \theta(b_i)y^{b_i} \rangle_{\theta \in M}}$$

where H_{orb}^* is the orbifold Cohomology ring, H^* the usual cohomology ring, $\langle . \rangle$ means the ideal generated by $.$, and $\mathbb{Q}[N]^{\Sigma^e}$ is $\oplus_{c \in N} y^c$ where y shows formal variables and the multiplication $y^c \cdot y^{c'} = y^{c+c'}$ when there is $\sigma \in \Sigma$ such that $\beta^e(c)$ and $\beta^e(c') \in \sigma$, and 0 otherwise, $[J]$.

(4) We apply the situation in (3) to the homogeneous manifold G/V of the principal bundle $V \hookrightarrow G \rightarrow G/V = D$ and the homogeneous vector bundle $\mathcal{E}$ associated to a representation $\rho : V \rightarrow Gl(H)$, for H a complex vector space. The formula in (11) translates into

$$(12) \quad H_{orb}^*([\mathcal{E} = G \times_V H]) = \frac{H^*(D_M)[\mathbf{m}]^\Sigma}{\langle \sum_i c_1(\psi_\lambda) + \lambda(b_i)y^{b_i} \rangle_\lambda}$$

The line bundles ψ_λ are coming from the action of the torus T , i.e the roots of the representation ρ . The equation (12) explains that the relations in characteristic cohomology can be given by polynomial equations in terms of first chern classes of line bundles associated to the co-characters of a maximal torus T in G . The same method can be applied to the principal bundle $P \hookrightarrow G_\mathbb{C} \rightarrow \check{D}$ where P is a parabolic subgroup. The curvature form Θ of a homogeneous manifold of this type can always be expressed as

$$\Theta = A \wedge^t \bar{A} - B \wedge^t \bar{B}$$

where A, B are matrices of $(1, 0)$ -forms. Using Cartan structure equation one obtains the following explicit form of Θ ,

$$\Theta = - \sum_{\alpha, \beta \in \Delta_+ - \Phi} [e_\alpha, e_\beta]_{\mathfrak{v}} \otimes \omega^\alpha \wedge \bar{\omega}^\beta$$

where Δ_+ is a system of positive roots for the root decomposition $\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta} \mathfrak{g}_\alpha$, $e_\alpha \in \mathfrak{h}^\vee$ the root vectors, $\omega^\alpha = e_\alpha^\vee$ and $\mathfrak{v} = \text{Lie}(V)$, Φ the corresponding sub-root system for V .

REFERENCES

- [FMN] B. Fantechi, E. Mann, F. Nironi, Smooth toric Deligne-Mumford Stacks, J. reine angew. Math. 648, 2010, 201-244
- [GGK] M. Green, P. Griffiths, M. Kerr, Mumford-Tate domains, Bollettino dell' UMI (9) III (2010), 281-307
- [J] Y. Jiang, The orbifold cohomology ring of simplicial toric stack bundles, Illinois J. of Math. Vol 52, No 2, 2008, 493-514
- [FMN] B. Fantechi, E. Mann, F. Nironi, Smooth toric Deligne-Mumford Stacks, J. reine angew. Math. 648, 2010, 201-244
- [GGK] M. Green, P. Griffiths, M. Kerr, Mumford-Tate domains, Bollettino dell' UMI (9) III (2010), 281-307
- [GS] A. Geraschenko, M. Satriano, Toric Stacks I, Theory of stacky fans, Trans. Amer. Mat. soc., Vol. 367, No 2, 2014, 1033-1071
- [KP] M. Kerr, G. Pearlstein, Boundary components of Mumford-Tate domains, arxiv preprint
- [KB] K. Behrend, Cohomology of Stacks, University of British Columbia, Lectures given at the school and conference on intersection theory and moduli, Sep. 9-27, 2002
- [TH] T. Hayama, On the boundary of moduli spaces of log Hodge structures, II: Non-trivial torsors, Nagoya Math. J. Volume 213 (2014), 1-20.
- [WF] W. Fulton, Introduction to toric varieties, princeton University press, Princeton, New Jersey 1993

,

E-mail address: mrahmati@cimat.mx, dr.khurramshabbir@gcu.edu.pk